

Contents of the New York Medical Abstract.

NOVEMBER, 1882.

The Spasmodic Paralyzes of Infancy, I., James Ross, M.D., F.R.C.P., 2 illustrations.
Hysterical Spine, Osman Vincent, F.R.C.S., Ed., M.R.C.S., Eng.
Case of Congenital Absence of Uterus and Vagina, William M. Reid, M.D., 1 illustration.
Two Cases of Pericarditis, with Remarks, Robert Saundby, M.D., Edin.
The Data of Alienism, I., Charles Mercier, M.B., Lond., F.R.C.S.
The Invasion of Inanimate Material by Granulation Tissue, C. B. Ball, M.D., F.R.C.S.I.
Puerperal Fever: Its Treatment and Prevention, John Lowe, L.R.C.S.I., &c.
A Few Words About Eatables, C. B. Radcliffe, M.D.
Two New Methods of Reduction in Dislocations of Femur, J. E. Kelly, F.R.C.S.I., 2 cuts.
Albuminuria of Epilepsy, Robert Saundby, M.D., Edin.
Pathology of Rickets, G. R. Gilruth, F.R.S.E.
Salicylate of Soda in Tonsillitis, and as a Local Application in Gout, Edward Mackey, M.D.
Observations on Chronic Morphinism, H. Obersteiner, M.D., Vienna.
Study of the Face as an Index of the Brain, Francis Warner, M.D., 6 illustrations.
Health of Criminal Women, F. M. Mosher, M.D.
Items.—Notes and Titles.

DECEMBER, 1882.

Four and a Half Years' Experience in the Lock Wards of the Edinburgh Royal Infirmary, A. G. Miller, F.R.C.S.E., 1 plate and 2 illustrations.
Localised Convulsions from Tumour of the Brain, J. Hughlings-Jackson, M.D., F.R.S., 2 illus.
Notes on Abdominal Surgery, William Stokes.
The Contagium of Gonorrhœa, C. R. Drysdale, M.D., M.R.C.P., Lond.
Offensive Catamenia or Bromo-Menorrhœa, Alfred Wiltshire, M.D., F.R.C., Lond.
Shearing or Moulding of Fœtal Head, in its Medico-Legal Aspect, W. J. Simpson, M.D., 2 plates.
Diagnosis of Cervical Tumours, Thomas E. Chavasse, M.D., C.M., F.R.C.S.
The Silica Spinal Jacket, George R. Fowler, M.D.
Chronic Dysentery, C. R. Francis, M.B., M.R.C.P.
Treatment of Diphtheritic Sore Throat, W. Allan Jamieson, M.D., F.R.C.P.
Somewhat Unusual and Complicated Case of Inguinal Hernia, Joseph Bell, F.R.C.S., Ed.
Clinical Lectures on Diseases of Women. No. I., J. Matthews Duncan, M.D., LL.D.
Indications for the Use of Digitalis, J. Milner Fothergill, M.D., Edin.
Items.—Notes and Titles.

MEDICAL ABSTRACT REPRINT.

No. 6.

RESECTION OF THE PYLORUS.

BY M. LE DOCTEUR KAHN.

JANUARY, 1883.

Intra-Peritoneal Pelvic Hæmatocele, Angus Macdonald, M.D.
Diabetic Coma, Balthazar Foster, M.D., F.R.C.P., and Robert Saundby, M.D., M.R.C.P.
The Germ Theory, Kendal Franks, M.D.
The Data of Alienism, II., Charles Mercier, M.B., Lond., F.R.C.S.
Some Points in Relation to Sounds of the Heart, W. H. Broadbent, M.D., F.R.C.S.
Clinical Lectures on Diseases of Women, II., J. Matthews Duncan, M.D., LL.D.
Whooping-Cough: Its Pathology and Treatment, Thomas M. Dolan, F.R.C.S., Edin.
Hysteria in Young Boys, M. Charcot, 4 illustrations, Translated from the French.
Items.—Notes and Titles.

FEBRUARY, 1883.

The Spasmodic Paralyzes of Infancy, II., James Ross, M.D., F.R.C.P., 2 illustrations.
Phthisis of the Larynx, G. Hunter MacKenzie, M.D., 1 plate, 2 illustrations.
Situation of Eggs and Fetuses in Twin Pregnancy; Symptoms, M. P. Buden, 8 illustrations.
Parasites in the Pork Supply of Montreal, Wm. Osler, M.D., and A. W. Clement.
Two Cases of Hysterical Contraction of Traumatic Origin, M. Charcot, Translated from the French, 2 illustrations.
Method for the Estimation of Urea in the Blood, John Berry Haycraft, M.B., F.R.S.E.
Therapeutic Value of Sulphurous Acid and Steel in Scarlatina Maligna, Keith Norman Macdonald, M.D.
Diseases of the Fallopian Tubes, Thomas Savage, M.D., M.R.C.P., Lond.
On the Pathology of General Paralysis, Joseph Wigglesworth, M.D., 1 plate.
Items.—Notes and Titles.

MARCH, 1883.

Two Cases of Hysterical Contraction of Traumatic Origin, M. Charcot, 5 illustrations, Translated from the French.
Koch's Researches on Tuberculosis, George A. Heron, M.D., Lond.
The Phenomena of Angina Pectoris, and their Bearing upon the Theory of Counter Irritation, W. Allen Sturge, M.D.
Treatment of Epilepsy, Robert Saundby, M.D., Edin.
Clinical Lecture on Diseases of the Lower Bowel, Edward Hamilton, F.R.C.S.I., 2 plates and 5 illustrations.
Washing, Medication and Alimentation of the Stomach, M. Dujardin Beaumetz, from the French
A Form of Loss of Memory Occasionally Following Cranial Injuries, Joseph Bell.
Nitrite of Sodium in Treatment of Angina Pectoris, Matthew Hay, M.D.
Peroxide of Hydrogen in Ophthalmic Practice.
Germs and the Spray, John Duncan.
Inoculation and Propagation of the Bacillus Tuberculosis, M. Babés, Paris, from the French.
Items.—Notes and Titles.

Office NEW YORK MEDICAL ABSTRACT,
93 Fulton St., New York

DEAR SIR:—This reprint is taken from the NEW YORK MEDICAL ABSTRACT, a journal with which you may not be acquainted, but which offers from month to month, in a condensed form, and at a very low price, some of the most mature and studied articles appearing in the

CURRENT MEDICAL LITERATURE OF EUROPE.

The ABSTRACT is now in the hands of many of our most prominent physicians, who have expressed their high appreciation of its value.

IT IS ALMOST AN ENCYCLOPÆDIA.

Its condensed form, its constant variety, the high merit and completeness of its articles, its many illustrations, and its low price, make it

ONE OF THE BEST AND CHEAPEST JOURNALS

in this country, and very valuable as a work of reference. To the busy practitioner who desires an intimate acquaintance with the most recent and best medical literature of Europe, the ABSTRACT is invaluable.

Specimen copy sent on application.

Yours very truly,

N. Y. MED. ABSTRACT.

ADVANTAGES OF THE NEW YORK MEDICAL ABSTRACT:

Its manner of condensing, by which *the language of the originals is retained*, while matter of local interest only is omitted. The completeness and high nature of its articles; valuable for present reading and future reference. Its condensed form and low price. Its many illustrations; no illustration appearing in the original being omitted on account of price.

In addition to the above we give, at the end of each number, the titles of the principal articles appearing in the current medical journals of this country and Great Britain during the previous month; and with the December number we publish a full *classified* index; together with a separate index under the authors' names.

Forty or more pages monthly, one dollar a year. Bound volumes, 1882, in cloth, \$1.50; half morocco, \$2.00.



Kahn. (Dr.)

Basiliysis for Dystocia from Hypertrophic Elongation of the Cervix Uteri, Alexander Russell Simpson, M.D., F.R.C.S., 3 illustrations.
So-Called Sponge-Grafting, Kendal Franks, M.D., F.R.C.S.I., 1 plate.
Chronic Acid in Certain Affections of the Tongue, Henry T. Butlin, F.R.C.S.
On the Character and Hallucinations of Joan of Arc, William W. Ireland, M.D.
The Prophylaxis of Ophthalmia Neonatorum, Alexander Russell Simpson, M.D., F.R.S.E.
Venesection, John Wylie, M.B.
Experimental Researches in the Employment of the Magnet in Epilepsy, MM. Bourneville and Bricon, from the French, 6 illustrations.
The Data of Alienism, III., Charles Mercier, M.B., Lond., F.R.C.S.
Clinical Remarks on Functional Vomiting of Hysteria, J. S. Bristowe, M.D., F.R.C.S.
Resection of the Pylorus, M. Le Docteur Kahn.
Items.—Notes and Titles.


MAY, 1883.

Two Cases of Tumour of the Larynx, D. N. Knox, M.A., M.B., 1 illustration.
Differential Diagnosis of Traumatic Lesions of the Brain, M. Duplay, *Faculté de Médecine*.
Clinical Lectures on Diseases of the Lower Bowel, Edward Hamilton, F.R.C.S.I.
Report to the Association for the Advancement of Medicine by Research on the Relation of Micro-Organisms to Tuberculosis, W. Watson Cheyne, M.B., F.R.C.S., 2 color-plates.
The Mental Condition in Hypnotism, D. Hack Tuke, F.R.C.P.
Items.—Notes and Titles.

RESECTION OF THE PYLORUS.

BY M. LE DOCTEUR KAHN.

Translated from *Le Progrès Médical*.



THE question of the resection of the pylorus is sufficiently old, for in the earlier years of this century a German physician, Karl Merrem, made such experiments upon a dog as demonstrated the possibility of the operation, and spoke of a Philadelphia surgeon who had made similar researches. Neither of them met with great success. People talked for a while of "Merrem's dream;" then the matter was so completely forgotten that when, in 1874, Gussenbauer and Winiwarter performed the same experiments they did not know of Merrem's labors in the year 1810. After several failures they succeeded in securing the survival of one of their animals—a dog—who was healthy and fat seven months after the operation, when he was sacrificed in experiments of another nature.

Kaiser and Werth then took up the question with the object of establishing more clearly the method of operation. In one case they even removed the entire stomach, drawing together the pylorus and cardia. The animal lived, and at the end of eight months weighed one half more than he did before the operation.

The operation upon the human subject was made for the first time on the 9th of April, 1879, by Péan, a Frenchman, upon a patient who threatened suicide if the surgeon refused to extirpate the cancer.

Eighteen months later, the 19th of November, 1880, Rydygier, of Kulm, operated upon a man of 64 years. The unsuccessful termination of these two operations (for the first patient died in five days, and the second in 12 hours), was not of a nature to attract many imitators; but Billroth, on the 22d of January, 1881, performed the operation upon a woman of 43 years, and she survived. This success made a great noise in Germany and excited much enthusiasm. Billroth hastened to operate upon two other cases, and his example was followed by many imitators. In the four months following, 9 resections of the pylorus were made by German physicians, and the year 1881 saw no less than 17 of them. To his first success Billroth had joined a second, and Wœlfler, his assistant, and Czerny added two more. Unfortunately Billroth's first patient died of relapse four months after the operation, and the absence of news from his second patient and the one operated upon by Czerny, left some doubt of the permanence of the attempted cure. It is known that all of the others died from the operation, and these facts explain why the enthusiasm concerning it has cooled.

But if Germany appeared calmed upon this point, if we do not find there any case of an operation made in 1882, we observe, nevertheless, that resection has since made the tour of the world, and we find it in Holland, England, Italy, Brazil, and the United States. We cannot present full details of the cases, but the following table gives the principal facts concerning the 27 cases of resection published up to the end of 1882.

Space will not permit us to give the complete manual of operation as thus far followed. We will draw attention to one of the points, which is interesting on account of the difficulty which it presents and the ingenious manner in which it

Operator and date of the Operation, Sex and Age.	Symptoms.	Nature of the Lesion.	Results.	Remarks and Bibliography.
1. J. Péan, At Paris, April 9, 1879. M.	Vomitings, Chæxia, very great dilatation of the stomach.	Cancer of the pylorus.	Two transfusions of blood. Death on the 5th day.	Péan Diag. et Trait. des Tum. de l'Abd. Gaz. des Hôpitaux, 1879, No. 60.
2. Rydygier, At Köln, Nov. 16, 1880. M., 64 years 9 mos.	Pains, vomitings, movable tumor above the umbilicus.	Scirrhus.	Died in 12 hours.	Dent. Zeitsch. für Chir., Bd. xiv.
3. Billroth, Vienna, Jan. 29, 1881. F., 43 years.	Vomitings, sudden appearance of symptoms in Oct. 1880. Tumor in region of pylorus.	Colloid.	Survived 4 months. Died May 23 of relapse, disease extending to walls of stomach and intestine.	Wolff. Ueber die von H. P. Billroth, ausg. ressect. der Carc. Pylorus, Vienna, 1881.
4. Billroth, Feb. 28, 1881. F., 39 years.	No vomitings. Tumor at left of umbilicus.	Epithelial.	Died in 9 days.	Daily vomitings after operation. A reopening on the 7th day and establishment of duodenal fistula. Wolff. Ueber die von H. P. Billroth, ausg. ressect. der carc. Pylorus, Vienna, 1881.
5. Billroth, March 12, 1881. F., 38 years.	Pains lasting one year. Vomitings. Stools very rare. Tumor above and right of umbilicus.	Medullary.	Died same evening.	Cancer adhered to pancreas. Wolff. Ueber die von H. P. Billroth, ausg. ressect. der carc. Pylorus, Vienna, 1881.
6. Nicolaysen, March 17, 1881. F., 37 years.	Pains for 10 months. Vomitings. Movable tumor. Dilatation of stomach.	Epithelial.	Died same night.	Some large ganglia removed from omentum. Nordström. Med. Arch. Bd. xii.
7. Bardenheuer, March 18, 1881. F., 54 years.		Cancer.	Died 26 hours after operation.	Vomitings of blood followed operation. At autopsy much blood found in stomach. Bardenheuer.
8. Wollner, April 8, 1881. F., 53 years.	Great dilatation of stomach. Tumor movable.	Colloid.	A year afterward patient still in very good health.	Wollner. Ueber die von H. P. Billroth, ausg. ressect. der Carc. Pylorus, Vienna, 1881.
9. Berns, April 23, 1881. F., 49 years.	Sick for 3 years. Dilatation of stomach. Tumor in umbilical region.	Cancer of pylorus.	Died in 4 hours.	Tumor very adherent to pancreas. During operation vena cava opened to a length of 10 centimetres. Dent. Zeitsch. für Chir., No. 50.
10. Jurié, May, 1881.	Tumor seemed to be very mobile.	Cancer of pylorus.	Speedy death.	Tumor, despite apparent mobility, adhered very strongly to pancreas. Dent. Zeitsch. für Chir., Bd. xvi.
11. Czerny, June 21, 1881. M., 28 years.	Pains and vomitings. Symptoms going back no further than six weeks.	Colloid.	In Jan., 1882, patient still very well, and had increased much in weight.	pa-Arch. de Langenbeck, Bd. 27, Hft. 4.
12. Lucke, June 25, 1881. M., 33 years.	Pains. No vomitings. Regular stools. Tumor painful. General condition good.	Scirrhus.	Died in 10 hours.	At autopsy heart soft, valves injected, lungs oedematous, scirrhus ganglia back of pancreas. Dent. Zeitsch. für Chir., Bd. xvi.
13. Kronlein, June 4, 1881. F., 54 years.	Pains for 3 months. Tumor very movable. Very marked emaciation.	Cancer.	Died in 24 hours.	Despite its mobility, tumor adhered very strongly. Carcinomatous infiltration extended into the liver and ganglia. Dent. Zeitsch. für Chir., Bd. xvi.
14. Kitajewsky, July 16, 1881. F., 52 years.	Very weak.	Cancer.	Died in 6 hours.	Sutures of oclusions found a little relaxed at autopsy. Dent. Zeitsch. für Chir., 1881, p. 783.

Operator and date of the operation. Sex and Age.	Symptoms.	Nature of the Lesion.	Results.	Remarks and Bibliography.
15. Weisslechner, Aug. 18, 1881. M., 47 years.	Very cachectic.	Cancer.	Died in 5 hours.	Tumor adhered to pancreas and liver. <i>Cent. f. Chir.</i> , 1881, p. 783.
16. Billroth, Oct. 23, 1881. F., 36 years.	Pains for 4 months. Vomiting. Small tumor.	Glandular.	Five months after-ward patient in very good health.	<i>Wien. Med. Woch.</i> , 1881, No. 51; 1882, No. 14.
17. Billroth, Nov. 5, 1881. M., 44 years.	Great dilatation of stomach.	Cancer.	Died on the third day.	Tumor adhered to pancreas. <i>Wien. Med. Woch.</i> , 1881, No. 51; 1882, No. 14.
18. Bardenheuer.		Cancer.	Died on the eighth day.	Rydygier. <i>Sammlung Kl. Fort.</i> , No. 220.
19. Bardenheuer.		Cancer.	Died on the second day.	Tumor adhered to pancreas. Rydygier. <i>Sammlung Kl. Fort.</i> , No. 220.
20. Langenbeck.			Died soon after-ward.	Strong adhesion to pancreas, which was deeply involved. Rydygier. <i>Sammlung</i> , etc., 11th Congress German Surgeons.
21. Gussenbauer, F.	Tumor very movable.		Died in 16 hours.	Extended adhesions to pancreas, of which it was necessary to remove a portion. Rydygier. <i>Sammlung</i> .
Operator and date of the operation. Sex and Age.	Symptoms.	Nature of the Lesion.	Results.	Remarks and Bibliography.
22. Southam, April 5, 1882. M., 43 years.	Pains, dilatation of stomach, vomitings, emaciation. Very pronounced constipation. Stools rare and tarry. Tumor movable.	Scirrhus.	Died in 14 hours.	<i>Brit. Med. Jour.</i> , 1882, No. 1, 126.
23. Fort, April 17, 1882. Rio Janeiro. F.	Extreme emaciation. Tumor movable.	Cancer.	Died very soon after operation.	Tumor adhered to pancreas, portal vein, and other portions of hepatic pecticle. <i>Gaz. des Hôp.</i> , 1882, No. 123.
24. Hahn, May 19, 1882. F., 63 years.	Pains, vomitings, emaciation, enormous dilatation of stomach. Tumor very mobile.	Cancer.	Died on the seventh day.	Patient had had fecaloid vomitings, which from the first led to a diagnosis of ileus. <i>Berlin Klin. Woch.</i> , 1882, No. 37.
25. Richter, May 25, 1882. M., 51 years.	Pains, vomitings, emaciation, tumor.	Cancer.	Died in 3 hours.	<i>San Francisco West. Lancet.</i> , July, 1882. Reported by Centralbl., 1882.
26. Caselli, June 14, 1882. F.	Dilatation, pains.	Cancer.	Died in 7 hours.	At the autopsy the other organs were found healthy. <i>Italia Medica</i> , June, 1882.
27. Kohler, F., 65 years.	Symptoms of cancer had been diagnosed 6 mos. previously.	Cancer.	Died in 6 hours.	<i>The Med. Herald</i> , 1882, No. 41. Reported by Centralbl., 1882.

is surmounted. We refer to that part of the operation which consists in securing the duodenum to the stomach.

In consequence of the frequent dilatation of the stomach, the calibre of its section is much larger than the duodenal, so that we have two openings of different diameters to unite. Péan, in his operation, found it very difficult to reduce the borders of the stomachal section sufficiently to adapt it to the duodenum. The German surgeons have discovered an ingenious method of doing it easily. Rydygier had, for this purpose, excised a triangular piece from the great curvature, sutured the edges of this section, and adapted it to the duodenum. Billroth, operating in a more simple way, folded in the excess of membrane and sutured it to the side of the stomach. But as the duodenal and stomachal sections had been made vertically, this disposition of the membrane brought about, in case number 4, the formation of a *cul de sac* at the extremity of the greater curve, which made the passage of food more difficult and caused more vomitings than the patient had experienced before the operation. He modified his method as follows: He divided the stomach and duodenum, not vertically as before, but following an oblique line from top to bottom and from left to right—thus making a larger opening of this portion than by cutting vertically—and sutured the stomachal section from the smaller bend. A third method has been proposed. It consists in cutting the duodenum vertically, and the stomach in a broken line composed of three parts, the superior to be oblique from top to bottom and from left to right from the smaller bend, and the inferior oblique from top to bottom and from right to left joining the greater curve. The median line is then cut, vertically and between the two, joining one to the other. The superior and inferior incisions are then sutured, the median serving to receive the duodenum.

In examining the results of the resection of the pylorus as thus far obtained, what do we find? Out of 27 persons operated upon, only four have survived; the 23 others died within a few hours or days after the operation. Of these 23, a few only showed signs of peritonitis, the greater part died in collapse. The mortality has been frightful, and it is proper to ask if an operation should be permitted which has thus far shown itself so murderous. Its partisans object, certainly, that we should not decide from the cases recorded in 1881-2 because of the bad condition of the patients who were operated upon, they for the most part presenting adhesions of the pylorus to the pancreas, a gland that they were obliged to cut into in order to remove all the parts involved by the disease. This latter is of course a matter which should always be the rule in an operation which has for its end the extirpation of cancer. They add that to-day they would not operate in such cases and that everyone now regards an adhesion to the pancreas as an absolute contra-indication.

In order that there may be no adhesion, it would be necessary to operate early, before the disease had time to spread. But at this time would the diagnosis be sufficiently well established to warrant an operation? We know the difficulties so often encountered in this disease well enough to reply that in most cases the diagnosis will only be established when it is too late.

But let us suppose the most favorable case, that is to say, one having an assured diagnosis and symptoms which have been manifested but a short time, conditions presented by four choice cases of which we shall speak further on. Have we in such subjects any means of knowing if the pylorus is still at liberty or if it is already adherent to the pancreas? To this question we must say, no!

In fact a sign that one would willingly take for a proof of the want of adhesion, is the perfect mobility of the tumor. In these 4 cases—Nos. 10, 12, 13 and 21—the mobility was perfect, and yet at the time of operation there were found very extended adhesions to the pancreas.

On the 25th of June, 1881, Lücke, of Strasbourg, operated upon a patient of 35 years who had never been sick before, had felt pains for only 5 months, had kept his appetite and had not been troubled with vomitings. The stools were regular and free from blood. The patient was anæmic but not cachectic. Here certainly was a group of symptoms which it would be difficult to improve upon. One might even ask how Lücke was able to decide upon risking an operation on a patient in so good a condition. Let us add that the tumor had an extreme mobility, yet at the time of operation it was found adhering to the pancreas and the patient died ten hours afterward.

[Continued in April ABSTRACT. Specimen copy sent on application.]

FOR PHYSICIAN'S USE ONLY.

SVAPNIA

OR

PURIFIED OPIUM.

CONTAINS THE
ANODYNE AND SOPORIFIC

ALKALOIDS

CODEIA,

NARCEIA

AND

MORPHIA.



EXCLUDES THE
POISONOUS & CONVULSIVE

ALKALOIDS

THEBAIN,

NARCOTIN

AND

PAPAVERRIN.

DOSE, THE SAME AS OPIUM.

This article is not intended for popular use, but only on prescription of the profession. It is to take the place of Opium in cases where that drug acts injuriously.

Dr. John Harley, of London, in his "Old Vegetable Neurotics," details a large number of experiments upon the human and animal system, with six of what he considers the narcotic alkaloids. He concludes that although all six possess both narcotic and hypnotic properties, yet these are so varied in degree and force, as to make their effects, when exhibited singly, very distinct from those following their exhibition in combination.

Taking the experience of practical physicians with Dr. Harley's results, as a basis, we would group them in the following order:

First Group.
Anodyne and Hypnotic Elements.

1. Morphia.
2. Narceia.
3. Codeia.

Second Group.
Narcotic and Convulsive Elements.

1. Thebain.
2. Cryptopin.
3. Papaverin.

Now SVAPNIA is a distinctive name given to the first group, representing the ANODYNE and HYPNOTIC ELEMENTS; the second group, or the narcotic and convulsive elements of Opium being eliminated; and is not, therefore, a simple principle, or a single constituent of Opium.

The relative values of each being known, we can select and utilize those that are valuable, and reject those known to be deleterious and inert.

This is what we claim, and all we claim for SVAPNIA. It can be relied upon and given in all cases where Opium or Morphia is indicated with equally good effects; and in addition to this, there will be found in the practice of every physician, cases occurring almost every day, in which idiosyncrasy and peculiar states and diseases of the brain debar us from the use of Opium and Morphia, but where Svapnia can be exhibited with the happiest results.

In SVAPNIA, there is retained all the Morphia and the greater part of the Codeia and Narceia, but combined with the native acids of Opium, meconic and thebolactic, in such a manner as to render those constituents soluble and active.

SVAPNIA IS UNIFORM IN ITS PROPORTIONS, and is prepared to conform to a standard of Opium representing ten per cent. of Morphia. This is done by a simple proportion of the neutral excipient, after determining the actual amount of the alkaloids obtained from the Opium used.

We may safely affirm that the danger of bad results is much diminished by a resort to that preparation of Opium, in which the poisonous elements are eliminated, and the anodyne elements in such a state of combination, as to reduce their toxic, and enhance their hygienic effects, such as has been proved to be the case with SVAPNIA.

SOLE AGENTS:

KIDDER & LAIRD,
83 John Street, New York.

FOR CONSUMPTION AND WASTING DISEASES.

HYDROLEINE

Has been proved of the highest value in CONSUMPTION and all
WASTING DISEASES, invariably producing IMMEDIATE
INCREASE IN FLESH AND WEIGHT.

FORMULA of HYDROLEINE.

Each dose of two teaspoonfuls, equal to 120 drops, contains:

Pure Cod Liver Oil.....	80 m. (drops.)	Soda.....	1-3 grains.
Distilled Water.....	35 "	Boric Acid.....	1-4 "
Soluble Pancreatin.....	5 grains.	Hyochohic Acid.....	1-20 "

DOSE—Two teaspoonfuls alone, or mixed with twice the quantity of soft water, to be taken thrice daily with meals.

The principles upon which this discovery is based have been described in a Treatise on "THE DIGESTION AND ASSIMILATION OF FATS IN THE HUMAN BODY," by H. C. BARTLETT, Ph.D., F.C.S., and the experiments which were made, together with cases illustrating the effect of Hydrated Oil in practice, are concisely stated in a Treatise on "CONSUMPTION AND WASTING DISEASES," by G. OVEREND DREWRY, M.D.

In these Treatises the Chemistry and Physiology of the Digestion of the Fats and Oils is made clear, not only by the description of a large number of experiments scientifically conducted, but by cases in which the deductions are most fully borne out by the results.

Copies of these Valuable Works will be sent free on application.

HYDRATED OIL, HYDROLEINE, WATER AND OIL.

HYDROLEINE is readily tolerated by the most delicate stomachs, even when the pure Oil or the most carefully prepared Emulsions are rejected. The Oil is so treated with pancreatin, soda, boric and hyochohic acids, that the process of digestion is partially effected before the organs of the patient are called upon to act upon it. Consequently it is readily assimilated. It will nourish and produce increase in weight in those cases where oils or fats, not so treated, are difficult or impossible to digest. In CONSUMPTION and other WASTING DISEASES, the most prominent symptom is *emaciation*, of which the first is the starvation of the fatty tissues of the body, including the brain and nerves. This tendency to emaciation and loss of weight is arrested by the regular use of HYDROLEINE, which may be discontinued when the usual average weight has been permanently regained.

The permanence and perfection of the emulsion, and the extreme solubility of the HYDRATED OIL, solely prepared and sold by us under the name of HYDROLEINE, is shown by its retaining its cream-like condition as long as the purest Cod-Liver Oil will retain its sweetness. Unlike the preparations mentioned, or simple Cod-Liver Oil, it produces no unpleasant eructation or sense of nausea, and should be taken in such very much smaller doses, according to the directions, as will insure its complete assimilation; this, at the same time, renders its use economical in the highest degree.

To brain-workers of all classes, Hydrated Oil is invaluable, supplying, as it does, the true brain food.

Economical in use—certain in result. Tonic—Digestive and Highly Nutritive.

NEW PRINCIPLE FOR THE FAT. ASSIMILATION.

KIDDER & LAIRD, Agents for the United States,

Retail Price, 1.00 per Bottle.

Depot, 83 John Street, New York.